

---

## Contents :

---

Letter from the Editor .....	<b>2</b>
Third-time recurrence of CS scar endometriosis after one cesarean section: A case report and Review of literature <i>Ahmed Sherif Abdel Hamid, Suzy AbdelAziz Abdel Hamid</i> .....	<b>3</b>
Distal vaginal atresia presented with apareunia with regular menses and not primary amenorrhea diagnosed by 3D ultrasound and managed by vaginal pull-through aided by 3D ultrasound: A case report and review of literature <i>Ahmed Sherif Abdel Hamid, Mounir Mohamed Fawzy el Hao, Suzy Abdelaziz Abdel Hamid</i> .....	<b>10</b>
The value of subcutaneous Fucidic acid instillation during elective cesarean delivery in prevention of surgical site infection <i>Asmaa I Ogila, Yossra Lasheen, Ahmed H Attia</i> .....	<b>17</b>
Association between Follicular Fluid Estradiol and Clinical Pregnancy Outcome in Intracytoplasmic Sperm Injection Cycles <i>Hussein M., Al-Kady M., Abdel Wahab H., Rushdy E., Refaat N., Dina Y. Mansour</i> .....	<b>25</b>
Role of Routine Hysteroscopy in Management of Women with Unexplained Infertility <i>Amr M. El Helaly, Khaled S. Mohamed, Hanan H. ElKhateeb, Dina Y. Mansour</i> .....	<b>40</b>
Azithromycin versus Erythromycin in Preterm Premature Rupture of Membranes: Mansoura Experience <i>Hamsa Gomaa Ramadan, Hesham Mahmoud Shalan, Abd Elhady Abd Elhady Zayed, Ahmed Abdelhamid El-Zayadi</i> .....	<b>49</b>

---

Low Dose Corticosteroids in Management of Hyperemesis Gravidarum at Mansoura University Hospital <i>Hanan Abd Almonem Ahmed, Maged El-shamy, Emad Ahmed Fayala, Mahmoud Mohamed Awad</i> .....	<b>60</b>
Uterocervical angle and cervical length as predictors for preterm birth in low-risk women <i>Mohamed K. Etman, Vivian A. Youssef, Mohamed S. Bakry, Sahar MY. El-Baradie</i> .....	<b>72</b>
A pilot study of The Efficacy of Using Chromium Salts in Reducing Hirsutism Scoring & BMI in Polycystic Ovary Syndrome Patients: Double Blinded Randomized Controlled Trial <i>Dina Yahia Aly Mansour, Heba Mohammed Eid, Marwa Saber Snosi</i> .....	<b>79</b>
Role of Activated Natural Killer Cells (CD3, CD56, CD16) in Repeated Implantation Failure in Women Undergoing IVF/ICSI Cycles <i>Noura El-Nassery, Abdel Maguid Ramzy, Amal Shohayeb1, AyatAllah A Nassef, Ahmed M Maged, Nancy M Ismael</i> .....	<b>89</b>
Effect of Luteal Estrogen Priming in Women with Poor Ovarian Response Undergoing IVF Using Antagonist Protocol <i>Noura El-Nassery, Amal Shohayeb, Mostafa Gamea, Ahmed M Maged, Mahmoud Soliman</i> .....	<b>99</b>
Single versus Double Frozen Embryo Transfer in PCOS Patients <i>Walid M Elnagar, Khaled F Helal</i> .....	<b>106</b>

---

## Letter from the Editor:

---

*Dear esteemed colleagues,*

Warm greetings

We welcome your comments as well as the scientific activity to be incorporated in the upcoming issues. Very important subjects are included in this issue. Not all cases of vaginal atresia cause primary amenorrhea but may present later with sexual problems. 3D rectal ultrasound is an excellent aid in diagnosis and treatment aid. Follicular fluid E2 concentration had fair predictive value in oocyte maturation, fertilization, embryo quality, chemical and clinical pregnancy. But it was an independent predictor of MII-grade oocytes production. Hysteroscopy (diagnostic, therapeutic) should be performed in women with unexplained infertility. Azithromycin can be used in place of erythromycin for the expectant management of PPROM. Azithromycin only benefits from its availability and reduced gastrointestinal side effects. Corticosteroids were highly effective in reducing the severity of vomiting in hyper emesis gravidarum patients after 48 hours from the start of the treatment using PUQE score and improving their QOL score compared with the standard treatment in the control group. The uterocervical angle is a potential novel screening tool for predicting preterm birth better than cervical length. Adding chromium picolinate to metformin in PCOS patients caused further improvement of Hirsutism score and regularity of menses, and further decrease in BMI. Estrogen priming did not increase the number of MII retrieved ova in POR. Double blastocyst transfer (DBT) even of good quality worsens the outcomes of ICSI for PCOS women. DBT is significantly associated with small birth weight, high incidence of preterm labour, need for operative delivery and NICU admission with subsequent reduction of live birth rate.

Best regards.

***Aboubakr Elnashar***

*MD*

*Chief Editor of EFSSJ*

*Prof. obs Gyn. Benha university, Egypt*

*elnashar53@hotmail.com*